

## Unforeseen consequences of the COVID-19 pandemic: Increased frequency of kite-string injuries in magnificent frigatebirds (*Fregata magnificens*) in Rio de Janeiro state, Brazil

DW Goldberg<sup>\*†</sup>, RET Vanstreels<sup>‡</sup>, MM Alcalá<sup>†</sup>, R Hurtado<sup>‡</sup>, PF Mantovani<sup>§</sup>, LST Cunha<sup>#</sup>, PP Serafini<sup>†</sup> and AF Barbosa<sup>¶</sup>

<sup>†</sup> Econservation Estudos e Projetos Ambientais, Rua José Alexandre Buaiz 300, Enseada do Suá, Vitória, ES, 29050-580, Brazil

<sup>‡</sup> Institute of Research and Rehabilitation of Marine Animals (IPRAM), Cariacica, ES, Brazil

<sup>§</sup> Síntese Serviços Veterinários, São Paulo, SP, Brazil

<sup>#</sup> Laboratório de Radioisótopos Eduardo Penna Franca, Instituto de Biofísica Carlos Chagas Filho, Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, RJ, Brazil

<sup>†</sup> Centro Nacional de Pesquisa e Conservação de Aves Silvestres (CEMAVE), Instituto Chico Mendes para a Conservação da Biodiversidade (ICMBio), Florianópolis, SC, Brazil

<sup>¶</sup> Coordenação de Licenciamento Ambiental de Exploração de Petróleo e Gás, Instituto Brasileiro do Meio Ambiente e dos Recursos Naturais Renováveis (IBAMA), Rio de Janeiro, RJ, Brazil

\* Contact for correspondence: daphwrobel@gmail.com

### Abstract

Kite flying is a popular hobby and sport for children and adults. Despite being illegal in Rio de Janeiro state, Brazil, the use of abrasive threads remains widespread and poses a health risk to both humans and animals. In this study, we analysed the records of 462 magnificent frigatebirds (*Fregata magnificens*) submitted to rehabilitation centres or found dead along the southern coast of Rio de Janeiro state from October 2016 to August 2020. Of these, 244 individuals (52.8%) presented wing lesions consistent with kite-string injury, which can have a critical impact on the ability of frigatebirds to fly and will ultimately cause their death. Even when veterinary care is provided, only a small proportion of the individuals (2%) will fully recover the ability to fly in order to be released back to the wild. In 2020, an atypical increase in the number of individuals with kite-string injuries (~1,200% increase compared to other years) was noted in the weeks following the suspension of school activities and commerce in response to the COVID-19 pandemic. The number of frigatebirds with kite-string injuries recorded in a given week was positively correlated with internet searches for kite-related terms, which also peaked during the quarantine period of the COVID-19 pandemic. This illustrates how pandemic events may aggravate existing human-wildlife conflicts, and how preparedness plans need to incorporate measures to help communities cope with boredom and isolation during quarantine in ways that do not negatively impact the welfare and conservation of wildlife.

**Keywords:** animal welfare, human-wildlife conflict, kite flying, manjha, pandemic, seabird

### Introduction

Kite flying is a popular hobby and sport for children and adults all over the world, and sometimes includes kite fighting, where the objective is to cut the opponent's kite string by means of kite lines coated with a mixture of either ground glass and glue (known as 'manjha' or 'cerol'), or quartz powder and aluminum oxide ('linha chilena') (Babu *et al* 2015). As a result of the use of abrasive threads, an increasing number of kite-string injuries have been reported in both humans and animals (Ladeira *et al* 2012; Roy & Shastri 2013; Muvalia *et al* 2019). In India, kite-flying festivals pose a major threat to raptors, vultures, pigeons and parrots, amongst others, including endangered species, such as the white-rumped vulture (*Gyps bengalensis*) (Roy & Shastri 2013; Babu

*et al* 2015). In Brazil, kite-string injury is also a frequent cause of admission of birds to rehabilitation centres, especially birds of prey (Joppert 2007; Brito 2017).

Although abrasive kite threads have been prohibited in Rio de Janeiro since July 2019 (state law no 8478), their use remains widespread. In the first semester of 2020, when school activities and commerce were suspended due to the COVID-19 pandemic, magnificent frigatebirds (*Fregata magnificens*) in Rio de Janeiro began experiencing kite-string injuries at unprecedented levels (Figure 1). In this study, we quantify the occurrence of kite-string injuries in frigatebirds along the southern coast of Rio de Janeiro state, and discuss the need to incorporate measures that help communities cope with boredom and isolation during quarantine in a manner that does not negatively impact the welfare and conservation of wildlife.





**Table 1 Summary of the number of magnificent frigatebirds (*Fregata magnificens*) recorded along the southern coast of Rio de Janeiro state (October 2016–August 2020).**

Injury	Juvenile	Adult female	Adult male	Unknown age/sex	Total
Confirmed kite-string injury	1 (1: 0)	10 (9: 1)	2 (2: 0)	0	<b>13 (12: 1)</b>
Suspected kite-string injury	47 (43: 4)	92 (88: 4)	92 (90: 2)	233 (2: 231)	<b>231 (221: 10)</b>
Other injury	33 (24: 9)	38 (33: 5)	40 (37: 3)	5 (2: 3)	<b>116 (97: 19)</b>
No trauma	26 (22: 4)	34 (33: 1)	39 (34: 5)	3 (2: 1)	<b>106 (90: 12)</b>
Undetermined (advanced decomposition)	25 (0: 25)	20 (0: 20)	18 (0: 18)	64 (30: 34)	<b>127 (0: 127)</b>
<b>Total</b>	<b>132 (90: 42)</b>	<b>194 (163: 31)</b>	<b>191 (163: 28)</b>	<b>72 (34: 38)</b>	<b>589 (420: 169)</b>

Results presented as 'all individuals (live individuals: dead individuals).'

**Table 2 Summary of the rehabilitation outcomes of magnificent frigatebirds (*Fregata magnificens*) rescued alive along the southern coast of Rio de Janeiro state (October 2016–August 2020).**

Injury	Spontaneous death	Euthanasia	Release	Total
Kite-string injury (confirmed or suspected)	80 (36%)	140 (62%)	5 (2%)	<b>225</b>
Other injury	43 (49%)	30 (34%)	14 (16%)	<b>87</b>
No trauma	40 (42%)	10 (11%)	45 (47%)	<b>95</b>
<b>Total</b>	<b>163 (40%)</b>	<b>180 (44%)</b>	<b>64 (16%)</b>	<b>407</b>

Results are presented as 'number of individuals (percentage).'

veterinary care, and the rehabilitation outcome (release, euthanasia, or spontaneous death) was recorded.

School and commerce in Rio de Janeiro were suspended on 16 and 24 March 2020, respectively; commerce re-opened on 27 June 2020, and school activities were still suspended as of 23 August 2020. Google Trends (<https://trends.google.com/>) was used to obtain a weekly index of internet searches by users in Rio de Janeiro state from 3 October 2016 to 23 August 2020 by averaging the results for three search terms: 'pipa' (Portuguese for 'kite'), 'cerol', and 'linha chilena.' Human population density at the locations where frigatebirds were found was derived from Global Population of the World v4 (Doxsey-Whitfield *et al* 2015). The geodesic kernel density of frigatebirds recorded by the beach-monitoring programme during the study period was obtained using the Kernel Density tool as implemented in ArcGIS 10 (ESRI, Redlands, USA), with a cell size of 0.001 km<sup>2</sup> and a search radius of 10,000 km. Data from an ongoing satellite tracking study are represented in this study for comparative purposes. The tracked individual was an adult female captured in August 2018 at Redonda Island, which was fitted with a Harrier-4 GPS-UHF tracker weighing 12–14 g (Ecotone Telemetry, Gdynia, Poland) using the backpack harness with Teflon tape; the data represented in this study were obtained from 1 to 30 September 2018.

Data normality was evaluated with the Kolmogorov-Smirnov test. Chi-squared tests were used to test whether

the categories of trauma were evenly distributed among the individual variables (context, status, developmental stage, and sex). The Kruskal-Wallis test with Bonferroni-corrected Mann-Whitney pair-wise tests were used to compare human population density among categories of trauma. Linear regression analysis was used to evaluate the relationship between the weekly number of frigatebirds with kite-string injuries and the index of kite-related internet searches.

## Results

Five hundred and eighty-nine magnificent frigatebirds were recorded during the study period (Table 1). Due to the advanced decomposition of some carcasses, only 462 individuals were evaluated for trauma, of which 244 (52.8%) had wing lesions consistent with kite-string injury (Figure 1). It is worth noting that 14 cases classified as 'Other injury' were the result of interaction with fishing gear (fishing lines or hooks). Records of frigatebirds were widely distributed throughout the study area, but individuals with kite-string injuries were predominantly concentrated along the densely populated margins of Guanabara Bay (Figure 2). Frigatebirds with kite-string injuries were found at locations with significantly higher human population density (median = 953 inhabitants km<sup>-2</sup>, n = 244) than frigatebirds with other injuries (296 inhabitants km<sup>-2</sup>, n = 116) or without trauma (226 inhabitants km<sup>-2</sup>, n = 102;  $H = 12.96$ ,  $df = 2$ ;  $P = 0.002$ ).

**Figure 3**

Time series of the number of magnificent frigatebirds (*Fregata magnificens*) with kite-string injuries (red line; suspected and confirmed cases combined; green line: only confirmed cases) and without kite-string injuries (blue line) relative to the index of kite-related internet searches (dashed black line), periods of school holidays or suspension (yellow shaded areas) and commerce closure due to the COVID-19 pandemic (pink shaded area) in Rio de Janeiro state, Brazil.

The occurrence of trauma was not unevenly distributed relative to the age group ( $n = 458$ ,  $\chi^2 = 4.329$ ,  $df = 2$ ;  $P = 0.115$ ), sex ( $n = 427$ ,  $\chi^2 = 0.048$ ,  $df = 2$ ;  $P = 0.976$ ), and context ( $n = 462$ ,  $\chi^2 = 5.064$ ,  $df = 2$ ;  $P = 0.079$ ). Kite-string injuries were more frequent in live (55%) than in dead individuals (26%) ( $n = 462$ ,  $\chi^2 = 14.539$ ,  $df = 2$ ;  $P < 0.001$ ). The rehabilitation outcome of live frigatebirds was markedly different depending on whether they had lesions consistent with kite-string injury ( $n = 407$ ,  $\chi^2 = 102.745$ ,  $df = 4$ ;  $P < 0.001$ ), with individuals with kite-string injuries experiencing similar levels of spontaneous death but being euthanased more frequently and released less frequently (Table 2).

From 2017–2019, the number of frigatebirds with kite-string injuries showed a tendency to increase around weeks 30–35 of each year, during or shortly after school winter holidays. In 2020, an atypical increase both in kite-related internet searches and in frigatebirds with kite-string injuries was noted between weeks 13–25 (29 March to 27 June), when commerce was suspended due to the COVID-19 pandemic (Figure 3). Sixty-one frigatebirds with kite-string injuries were recorded during this period, representing a 1,207% increase compared to the same period in 2017–2019 (mean = 4.7 individuals, range: 2–6). There was a significant positive correlation between the weekly number of frigatebirds with kite-string injuries and the index of kite-related internet searches ( $R^2 = 0.362$ ;  $P < 0.001$ ,  $n = 203$  weeks).

## Discussion

Frigatebirds are gliding seabirds that have the lowest body mass to wingspan ratio of all birds, allowing them to soar at high altitudes efficiently on weak ocean thermals (Weimerskirch *et al* 2003). Despite feeding primarily on marine prey, frigatebirds do not dive nor float on water; instead, they capture prey while gliding near the sea surface or harass other seabirds to force them to regurgitate and then steal their meal (Schreiber & Burger 2001). As such, these birds are reliant on the integrity of their wings in order to get their food. When frigatebirds injure their wings they may fall into the sea and drown; if they survive the fall, they may later starve due to an inability to fly. Additionally, the particularly fragile anatomy of their wings (narrow propatagial membrane, thin skin and hollow bones) render frigatebirds exceptionally susceptible to kite-string injuries.

Compared to other bird species (Roy & Shastri 2013; Babu *et al* 2015), injuries in this study were characterised by deep lacerations and muscle avulsion, with extensive damage to tendons, bones, nerves and blood vessels. Kite-string injuries that require tenorrhaphy (suturing of tendons) will often require a long period of post-surgical immobilisation and physical therapy, since tendons are poorly vascularised and heal slowly. In our experience, even when veterinary care is provided, only a small proportion of the individuals will recover the ability to fly (in this study, only 2% of the frigatebirds admitted with kite-string injuries could be released). Due to the unfeasibility of providing a sufficiently large permanent enclosure that would allow a frigatebird to take flight, individuals that are not fit for release after rehabilitation are euthanased.

Kite-injured frigatebirds were most frequently recorded in areas of high human population density, especially along the margins of Guanabara Bay. The frigatebird population from Cagarra and Redonda Islands are known to forage in large numbers at Guanabara and Sepetiba Bays, often flying over the city of Rio de Janeiro to cross from Guanabara Bay to Sepetiba Bay and *vice versa* (Figure 2[D]; L Cunha, personal observation 2020). Our results suggest that kites with abrasive threads might become a key cause of mortality for this population. As our data are derived from a Beach-Monitoring Project, with a monitoring effort that emphasises coastal environments, it is likely that the geographic distribution observed in this study underrepresents the occurrence of injured frigatebirds due to lower monitoring effort at inland environments (eg individuals that suffered severe injuries while flying over Rio de Janeiro city and which immediately fell in urban areas) and at coastal islands (eg individuals that suffered relatively mild injuries and were able to fly to seek refuge at their usual roosting/breeding sites). Moreover, because a proportion of frigatebirds injured by kite strings could have drowned and never washed ashore, this study may have underestimated the total number or the geographical distribution of kite-string injured individuals.

Kite flying is a popular sport during the winter holidays in Rio de Janeiro state, whereas beach activities tend to be

more popular during summer holidays (D Goldberg, personal observation 2020). This might explain why an increase in kite-injured frigatebirds is usually noted during and shortly after the school winter holidays. From February to June 2020, however, the number of kite-string injuries increased acutely (*circa* 1,200% compared to the same period in previous years). With the suspension of school activities on 16 March and the closure of commerce on 24 March due to the COVID-19 pandemic, many children and adults had plenty of free time to fly kites from their roof-tops, as this was one of the few leisure activities that was still safe during quarantine, especially in economically deprived communities. This was reflected in the number of internet searches using kite-related terms and, unfortunately, in the number of frigatebirds with kite-string injuries. The increase in the number of injured frigatebirds during the peak of the breeding season (May to August; Cunha 2018) at Redonda Island may represent an even more dramatic situation, as the injured adults will not return to the nest to protect or feed the chick, potentially resulting in increased chick mortality.

This study therefore illustrates how epidemic/pandemic events can have unforeseen consequences and aggravate pre-existing human-wildlife conflicts. This is not dissimilar to the impacts of wars and civil strife, which may amplify and create new direct and indirect impacts on wildlife through a multitude of mechanisms (Dudley *et al* 2002; Gaynor *et al* 2016). Since COVID-19 is the largest fast-spreading pandemic in the contemporary era, we still have limited understanding as to how such global events (and the measures employed to mitigate them) can lead to changes to institutional and socioeconomic factors that ultimately affect wildlife and nature conservation. Understanding these mechanisms at the regional level, taking into account wildlife communities, cultural and socioeconomic factors, will be critical in order to adequately prepare for and mitigate their impacts. Past experience has shown that strong and flexible partnerships between local communities, non-governmental organisations and international institutions are critical to mitigate the impacts of violent conflicts on wildlife (Dudley *et al* 2002; Gaynor *et al* 2016), and a similar approach might be necessary to deal with epidemic/pandemic events.

In the case of kite-string injuries in Brazil, in addition to a stricter enforcement of the prohibition of abrasive threads (especially for 'linha chilena', which cannot be easily home-made and is commercialised locally), other approaches beyond law enforcement are also clearly necessary. The development of a regional campaign that embraces kite flying as a positive leisure activity during quarantine, acknowledging its strong cultural roots in Brazilian culture and its role in promoting bonding between parents and their children, could be instrumental in raising awareness about the negative effects of abrasive threads to both humans and wildlife.

Since social isolation and quarantining can cause significant psychological stress (Reynolds *et al* 2008; Brooks *et al*

2020), the preparedness plans for epidemic/pandemic events would also benefit from planning on providing the population with safe leisure options. In the Rio de Janeiro context, for example, the distribution of kites with non-abrasive strings and other toys (eg action figures, spinning tops, marble balls, etc), perhaps with themes related to frigatebirds and other native species and accompanied by educational booklets and brochures, could help mitigate the social anxiety in ways that are not harmful to wildlife.

Furthermore, while frigatebirds are not the only birds affected by kite-string injuries in Rio de Janeiro, the fact that they are easily recognisable in flight and have several interesting characteristics (eg colourful gular pouch in males, ability to soar with great efficiency) makes them good candidates to become a flagship species. Awareness campaigns could therefore benefit from highlighting the beauty and value of this species, appealing to the public's empathy by showing the suffering experienced by injured frigatebirds and sharing uplifting stories of individuals that were successfully rehabilitated.

The relatively strong correlation between internet searches for kite-related terms and the occurrence of kite-string injuries in frigatebirds suggests that the internet might also be a potential vehicle for educational and enforcement activities. Where possible, the publication of information on how to manufacture 'cerol' or promoting the use of 'linha chilena' should be discouraged (eg demonetised or banned), and websites and online communities promoting kite flying must actively discourage the use of abrasive threads. Additionally, social media campaigns promoting safe kite-flying practices during strategic periods (eg school winter holidays and during quarantine periods) could also help.

Although magnificent frigatebirds are not currently considered threatened with extinction, the relatively high frequency of kite-string injuries documented in this study (244 individuals from October 2016 to August 2020) may be considered a significant threat at the regional level (breeding population = *circa* 3,400 pairs; Moraes-Ornellas & Ornellas 2012; Cunha *et al* 2013). Furthermore, it is worth noting that other species of frigatebirds are critically threatened, such as the Christmas frigatebird (*Fregata andrewsi*), which has a total population of *circa* 4,800 pairs (Birdlife International 2018) and the Atlantic lesser frigatebird (*Fregata trinitatis*), which has an estimated population of *circa* 50 individuals (Olson 2017). Considering the anatomical and behavioural similarities between these species, it seems likely that they would also be highly vulnerable to kite-string injury.

#### Animal welfare implications

Although the kite-string injuries are usually relatively small and do not damage vital organs, they have a critical impact on the frigatebirds' ability to fly and will ultimately cause their death due to drowning or starvation. The sudden increase in the number of magnificent frigatebirds with kite-string injury in Rio de Janeiro during the period of commerce closure due to quarantining in response to the COVID-19 pandemic illustrates how epidemic events may

aggravate existing human-wildlife conflicts. This highlights how pandemic events can have unexpected and far-reaching impacts on animal welfare, even when the pathogen does not infect animals. Pre-identifying human-wildlife conflicts will be valuable to inform preparedness plans for epidemic/pandemic events, so that these plans can incorporate measures to help communities cope with boredom and isolation during quarantine in ways that do not negatively impact the welfare and conservation of wildlife.

#### Acknowledgements

The Santos Basin Beach Monitoring Project (Projeto de Monitoramento de Praias da Bacia de Santos - PMP-BS) is one of the monitoring programmes required by Brazil's federal environmental agency, IBAMA, for the environmental licensing process of oil production and transport by Petrobras at the pre-salt pole (25°05'S 42°35'W to 25°55'S 43°34'W), between 2,100 m and 2,300 m isobaths. We are grateful to past and present PMP-BS teams, Petrobras, Instituto Brasileiro do Meio Ambiente e dos Recursos Naturais Renováveis (IBAMA), Instituto Chico Mendes para a Conservação da Biodiversidade (ICMBio) and WWF, for financing the satellite tracking study with the frigatebirds.

#### References

- Alves VS, Soares AB and Couto GS** 2004 Aves marinhas e aquáticas das ilhas do litoral do Estado do Rio de Janeiro. In: Branco JO (ed) *Aves Marinhas e Insulares Brasileiras: Bioecologia e Conservação* pp 83-100. Editora da Univali: Itajaí, Brazil. [Title translation: Marine and aquatic birds of the coastal islands of Rio de Janeiro state; Marine and insular birds of Brazil: Bioecology and Conservation]
- Babu S, Subramanya S and Dilawar M** 2015 Kite flying: Effect of Chinese manja on birds in Bangalore, India. *Indian Birds* 10: 13-18
- Birdlife International** 2018 *Fregata andrewsi*. *The IUCN Red List of Threatened Species*. <https://doi.org/10.2305/IUCN.UK.2018-2.RLTS.T22697742A132599384.en>
- Brito MCR** 2017 *Diagnóstico, registro e destinação da avifauna recebida no Pró-Arara Centro de Reabilitação de Animais Silvestres, Araras-SP*. Masters Thesis, Universidade Federal de São Carlos, Araras, Brazil. [Title translation: Diagnosis, records and rehabilitation outcomes of avifauna admitted at the Pró-Arara Wildlife Rehabilitation Center, Araras-SP]
- Brooks SK, Webster RK, Smith LE, Woodland L, Wessely S, Greenberg N and Rubin GJ** 2020 The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *The Lancet* 395: 915-920. [https://doi.org/10.1016/S0140-6736\(20\)30460-8](https://doi.org/10.1016/S0140-6736(20)30460-8)
- Cunha L** 2018 Seabirds. In: Bertoncini A, Moraes F, Borgonha M, Aguiar A and Duarte B (eds) *Marine Biodiversity and Diving Guide to the Islands of Rio* pp 313-323. Museu Nacional: Rio de Janeiro, Brazil
- Cunha LST, Alves VS, Rajão HB and Lanna AM** 2013 Aves do monumento natural das Ilhas Cagarras. In: Moraes F, Bertoncini A and Aguiar A (eds) *História, Pesquisa e Biodiversidade do Monumento Natural das Ilhas Cagarras* pp 177-205. Museu Nacional: Rio de Janeiro, Brazil. [Title translation: Birds of the Cagarras Islands Natural Monument; History, Research and Biodiversity of the Cagarras Islands Natural Monument]

- Doxsey-Whitfield E, MacManus K, Adamo SB, Pistolesi L, Squires J, Borkovska O and Baptista SR** 2015 Taking advantage of the improved availability of census data: a first look at the Gridded Population of the World, version 4. *Papers in Applied Geography 1*: 226-234. <https://doi.org/10.1080/23754931.2015.1014272>
- Dudley JP, Ginsberg JR, Plumptre AJ, Hart JA and Campos LC** 2002 Effects of war and civil strife on wildlife and wildlife habitats. *Conservation Biology 16*: 319-329. <https://doi.org/10.1046/j.1523-1739.2002.00306.x>
- Gaynor KM, Fiorella KJ, Gregory GH, Kurz DJ, Seto KL, Withey LS and Brashares JS** 2016 War and wildlife: linking armed conflict to conservation. *Frontiers in Ecology and the Environment 14*: 533-542. <https://doi.org/10.1002/fee.1433>
- Joppert AM** 2007 *Estudo prospectivo das causas de morte de Falconiformes e Strigiformes de vida livre no município de São Paulo*. PhD Thesis, University of São Paulo, São Paulo, Brazil. [Title translation: Prospective study of the cause of death of free-ranging Falconiformes and Strigiformes at São Paulo municipality]
- Ladeira RM, Carreiro PRL, Rezende-Neto JB, Iannuzzi GC and Elias AA** 2012 Epidemiology of powdered glass-coated kite line injuries: victims admitted to a trauma hospital in Belo Horizonte, Brazil. *Revista Brasileira de Epidemiologia 15*: 407-414. <https://doi.org/10.1590/S1415-790X2012000200018>
- Moraes-Ornellas VS and Ornellas RB** 2012 Notas sobre a colônia de aves marinhas da Ilha Jorge Grego - RJ. *Atualidades Ornitológicas 169*: 7-9. [Title translation: Notes about the seabird colonies at Jorge Grego island - RJ]
- Muvalia G, Jamshed N, Sinha TP and Bhoi S** 2019 Kite-string injuries: A case series. *International Journal of Critical Illness and Injury Science 9*: 147. [https://doi.org/10.4103/IJCIIS.IJCIIS\\_44\\_19](https://doi.org/10.4103/IJCIIS.IJCIIS_44_19)
- Olson SL** 2017 Species rank for the critically endangered Atlantic Lesser Frigatebird (*Fregata trinitatis*). *Wilson Journal of Ornithology 129*: 661-675. <https://doi.org/10.1676/1559-4491-129.4.661>
- Reynolds DL, Garay J, Deamond S, Moran MK, Gold W and Styra R** 2008 Understanding, compliance and psychological impact of the SARS quarantine experience. *Epidemiology & Infection 136*: 997-1007. <https://doi.org/10.1017/S0950268807009156>
- Roy A and Shastri K** 2013 Impact of kite string injuries and temporal variation in types of injuries and illnesses of white-rumped vultures of central Gujarat, India. *Journal of Threatened Taxa 5*: 4887-4892. <https://doi.org/10.11609/JOTT.o3451.4887-92>
- Schreiber EA and Burger J** 2001 *Biology of Marine Birds*. CRC Press: Boca Raton, Brazil. <https://doi.org/10.1201/9781420036305>
- Weimerskirch H, Chastel O, Barbraud C and Tostain O** 2003 Frigatebirds ride high on thermals. *Nature 421*: 333-334. <https://doi.org/10.1038/421333a>